

IBUPROFEN

eye-bew-pro-fen

What does it do?

Ibuprofen is an anti-inflammatory medicine used to reduce pain, inflammation and fever.

Before you start

- Tell your doctor if you have heart, kidney, liver, stomach or bowel problems, high blood pressure, asthma, or if you have ever had a stroke or TIA (mini-stroke).
- Tell your doctor if you have had problems with aspirin or other anti-inflammatory medicines.
- Tell your doctor if you are pregnant, planning to become pregnant, or breastfeeding.

How should you take it?

Take *ibuprofen* as directed.

Tablet or capsule: Take with a large glass of water.

Slow-release tablet: Take with a large glass of water. Swallow whole - do not crush or chew.

Liquid: Shake well before use. Measure each dose carefully with an oral syringe or measuring spoon.

What if you forget a dose?

If you take *ibuprofen* regularly and it is nearly time for your next dose, skip the missed dose and take your next dose at the usual time. Otherwise, take the missed dose as soon as you remember. Do not take two doses at the same time.

Can you take other medicines?

Some medicines available without a prescription may react with *ibuprofen* including:

- anti-inflammatories such as diclofenac (e.g. Voltaren Rapid®), or aspirin (e.g. Disprin®, in doses used for pain relief). These can also be found in some cold and flu medicines (e.g. Nurofen Cold and Flu®), and creams or gels (e.g. Voltaren Emulgel®)
- low-dose aspirin (e.g. Cartia®)

Tell your pharmacist or doctor about all medicines or treatments that you may be taking, including vitamins, herbal products or recreational drugs.

What side effects might you notice?

Side Effects	Recommended action
Symptoms of allergy including: skin rash, itching, swelling, trouble breathing Symptoms of a peptic ulcer including: tummy pain, red or black bowel motions, vomit that looks like coffee grounds Symptoms of liver problems including: yellow skin or eyes, itching, dark urine, pale bowel motions, abdominal pain	Tell your doctor immediately
Swollen feet or legs, short of breath	
Indigestion, nausea, diarrhoea	Take with food and tell your doctor if symptoms continue

If you notice any other effects, discuss them with your doctor or pharmacist.

Other information:

- It is safe to use paracetamol with *ibuprofen* for pain relief if you need to.
- Anti-inflammatory medicines can sometimes cause kidney problems, especially in people who are dehydrated. Don't take *ibuprofen* if you are dehydrated.
- Long-term use of anti-inflammatory medicines may increase the chances of a heart attack or stroke – talk to your health professional about your risk