



DABIGATRAN

da-bi-gat-ran

What does it do?

Dabigatran is used to prevent clots in your blood. It reduces your risk of having a stroke and stops new clots in your legs or lungs.

Before you start

- Tell your doctor if you have kidney, bleeding, liver or stomach problems.
- Tell your doctor if you are pregnant, planning to become pregnant, or breastfeeding.

How should you take it?

Take *dabigatran* regularly as directed.

Dabigatran can damage the oesophagus (food pipe). To avoid this, take it with a large glass of water. Swallow the capsule whole - do not crush, chew or open it. Sit or stand upright for at least 30 minutes after taking a dose.

What if you forget a dose?

If you take *dabigatran* twice a day:

If the next dose is less than six hours away, skip the missed dose and carry on as normal. If there are more than six hours until the next dose, take the missed dose as soon as you remember. Do not take two doses at the same time.

If you take *dabigatran* once a day:

Skip the missed dose and carry on as normal at the same time the next day. Do not take two doses at the same time.

Can you take other medicines?

Some medicines available without a prescription may react with *dabigatran* including:

- anti-inflammatories, such as diclofenac (e.g. Voltaren®), ibuprofen (e.g. Nurofen®), or aspirin (e.g. Disprin®, in doses used for pain relief). These can also be found in some cold and flu medicines (e.g. Nurofen Cold and Flu®).
- low-dose aspirin (e.g. Cartia®) - sometimes used with *dabigatran*, check with your doctor.

Tell your pharmacist or doctor about all medicines or treatments that you may be taking, including vitamins, herbal products (e.g. St John's wort) or recreational drugs.

What side effects might you notice?

| Side Effects | Recommended action |
|--|---|
| Reduced number of blood cells that help your blood to clot - symptoms include: easy or unusual bruising or bleeding Coughing or vomiting of blood, vomit that looks like coffee grounds Nose bleeds Red or dark brown urine, red or black bowel motions Chest pain, trouble breathing Headache, dizziness, changes in vision or speech Trouble swallowing, severe indigestion or heartburn, stomach pain | Tell your doctor immediately |
| Tiredness, pale skin | Tell your doctor |
| Stomach upset, heartburn, indigestion | Tell your doctor. Try taking with food. |

If you notice any other effects, discuss them with your doctor or pharmacist.

Other information:

- It is important to tell anyone who gives you medical or dental treatment (e.g. doctor, dentist, pharmacist or podiatrist) that you are taking *dabigatran*.