

ROPINIROLE

roe-pin-ih-role

What does it do?

Ropinirole is used to treat Parkinson's disease and restless legs syndrome. It acts like a chemical in your brain called dopamine. Dopamine helps your brain control your body movements.

Before you start

- Tell your doctor if you have low blood pressure, or mental health problems.
- Tell your doctor if you are pregnant, planning to become pregnant, or breastfeeding.

How should you take it?

Take *ropinirole* regularly as directed with a glass of water. You can take it with or without food.

Do not stop taking *ropinirole* suddenly without talking to your doctor first. Your doctor may reduce the dose gradually.

What if you forget a dose?

If it is nearly time for your next dose, skip the missed dose and take your next dose at the usual time. Otherwise, take the missed dose as soon as you remember. Do not take two doses at the same time.

Can you take other medicines?

Tell your pharmacist or doctor about all medicines or treatments that you may be taking, including vitamins, herbal products (e.g. kava) or recreational drugs.

What side effects might you notice?

Side Effects	Recommended action
Confusion, unusual behaviour or thinking Seeing, hearing or feeling things that aren't really there Unusual urges (e.g. gambling, eating, spending, sex) Fainting Suddenly falling asleep Squirming or wriggling movement (such as head wobble)	Tell your doctor
Abdominal pain, nausea, vomiting, constipation Dizziness, drowsiness, tiredness or weakness	Tell your doctor if troublesome
Lightheaded or dizzy after standing up	Stand up slowly. If it continues, or is severe, tell your doctor

If you notice any other effects, discuss them with your doctor or pharmacist.

Other information:

- *Ropinirole* can impair your ability to do tasks such as driving or using machines. Alcohol makes this worse. Discuss your risk with your health professional. See [Are you safe to drive?](#)
- Smoking can change the effect of *ropinirole*. Tell your doctor if you give up, cut down or start smoking.
- If you are taking *ropinirole* for restless leg syndrome, your symptoms might get worse when you start. Discuss this with your doctor, as changing the dose can help.