



DOMPERIDONE

dom-peer-ih-dohn

What does it do?

Domperidone is used to treat and prevent nausea and vomiting. It is also sometimes used for other conditions.

How should you take it?

Take *domperidone* as directed with a glass of water.

What if you forget a dose?

Should an occasional dose be missed it need not be taken later.

Can you take other medicines?

Some medicines available without a prescription may react with *domperidone* including:

- fluconazole (e.g. Diflucan®)
- antacids (e.g. Mylanta®) - do not take these within two hours of taking *domperidone*.

Tell your pharmacist or doctor about all medicines or treatments that you may be taking, including vitamins, herbal products (e.g. St John's wort) or recreational drugs.

What side effects might you notice?

Side Effects	Recommended action
Fast or irregular heartbeat, fainting	Tell your doctor immediately
Headache Dry mouth or throat Diarrhoea Sore or enlarged breasts, breastmilk production Irregular periods	Tell your doctor if troublesome

If you notice any other effects, discuss them with your doctor or pharmacist.

Other information:

- Tell your doctor if you have heart, bowel, liver or kidney problems, or if you have a pituitary tumour.
- Tell your doctor if you are pregnant, planning to become pregnant, or breastfeeding.
- Your doctor may do a heart test (ECG) before you start and while you are taking *domperidone*.
- Grapefruit, grapefruit juice or sour/Seville oranges may react with *domperidone*. Discuss with your pharmacist.