



ETHOSUXIMIDE

eth-oh-sux-ih-mide

What does it do?

Ethosuximide is used to treat and prevent seizures.

Before you start

- Tell your doctor if you are pregnant, planning to become pregnant, or breastfeeding.

How should you take it?

Take *ethosuximide* regularly as directed. You can take it with or without food.

Capsule: Take with a glass of water.

Liquid: Measure carefully with an oral syringe or measuring spoon.

What if you forget a dose?

Take the missed dose as soon as possible and continue as directed.

Can you take other medicines?

Some medicines available without a prescription may react with *ethosuximide* including:

- some antihistamines, such as chlorphenamine, hyoscine, meclozine and promethazine. These can be in medicines for allergies, nausea and colds.

Tell your pharmacist or doctor about all medicines or treatments that you may be taking, including vitamins, herbal products (e.g. St John's wort, ginkgo) or recreational drugs (e.g. ecstasy).

What side effects might you notice?

| Side Effects | Recommended action |
|---|---|
| Skin rash, skin peeling or blisters | Stop taking and see your doctor immediately |
| Suicidal thoughts Reduced number of blood cells that fight infections or help your blood to clot - symptoms include: fever, chills, sore throat or generally feeling unwell, or easy or unusual bruising or bleeding | Tell your doctor immediately |
| Loss of co-ordination/walking or handwriting problems Mood changes | Tell your doctor |
| Drowsiness, dizziness, headache Loss of appetite, weight loss Stomach upset | Tell your doctor if troublesome |

If you notice any other effects, discuss them with your doctor or pharmacist.

Other information:

- *Ethosuximide* can impair your ability to do tasks such as driving or using machines. Alcohol makes this worse. Discuss your risk with your health professional. (search NZTA - Are you safe to drive?)
- Do not stop taking *ethosuximide* without talking to your doctor first, unless you have a skin rash (see Side Effects).