



## OXYCODONE (long acting)

*ox-ee-koe-dohn*

### What does it do?

*Oxycodone* is used to relieve strong pain.

### Before you start

- Tell your doctor if you have bowel problems, sleep apnoea, if you have ever had a seizure, or have had a recent head injury.
- Tell your doctor if you are pregnant, planning to become pregnant, or breastfeeding.

### How should you take it?

Take *oxycodone* regularly as directed. You can take it with or without food. Swallow whole – do not crush or chew.

### What if you forget a dose?

If it is nearly time for your next dose, skip the missed dose and take your next dose at the usual time. Otherwise, take the missed dose as soon as you remember. Do not take two doses at the same time.

### Can you take other medicines?

Some medicines available without a prescription may react with *oxycodone* including:

- sedating antihistamines (e.g. Phenergan®)
- cough suppressants (e.g. Duro-Tuss®, Benadryl Dry Forte®)

Tell your pharmacist or doctor about all medicines or treatments that you may be taking, including vitamins, herbal products (e.g. St John's wort) or recreational drugs.

### What side effects might you notice?

Side Effects	Recommended action
Slow or shallow breathing, hard to wake up Seizures	Tell your doctor immediately
Confusion, hallucinations, mood changes Tummy pain (new or getting worse)	Tell your doctor
Constipation	Take your prescribed laxatives. Tell your doctor
Drowsiness, dizziness Nausea, vomiting, dry mouth or throat Itching, flushing, sweating Strange or uncontrolled movements	Tell your doctor if troublesome

If you notice any other effects, discuss them with your doctor or pharmacist.

### Other information:

- Take *oxycodone* 'long acting' regularly. If *oxycodone* is not relieving your pain contact your health professional - your dose may need changing.
- *Oxycodone* 'short acting' may be used with *oxycodone* 'long acting' for breakthrough pain.
- *Oxycodone* can impair your ability to do tasks such as driving or using machines. Alcohol makes this worse. Discuss your risk with your health professional. (search NZTA - Are you safe to drive?)
- *Oxycodone* may be addictive with long-term use.
- You may see the leftover shell of the tablet in your bowel motions – this is harmless.
- Long-term use of *oxycodone* may cause side effects such as less interest or trouble having sex, or changes in periods. It may also make some types of pain worse. Discuss with your doctor.
- If you have been taking *oxycodone* regularly for a long time, talk to your doctor before stopping.