



SUMATRIPTAN

soo-ma-trip-tan

What does it do?

Sumatriptan is used to treat migraines and cluster headaches.

Before you start

- Tell your health professional if you have heart or blood vessel problems, high blood pressure, or if you have ever had a stroke, or 'mini-stroke'.
- Tell your health professional if you are pregnant, planning to become pregnant, or breastfeeding.

How should you take it?

Take *sumatriptan* as directed when your symptoms first start.

Take the tablets with a large glass of water.

Inject the injection under the skin as directed.

If your symptoms get better, but then return, you can take another dose after 1 hour (injection) or 2 hours (tablet).

If *sumatriptan* does not help, do not take any further doses for the same attack, as it is unlikely to work. You can still use *sumatriptan* for a new attack.

Do not take more than the maximum daily dose advised by your health professional.

Can you take other medicines?

Some medicines available without a prescription may react with *sumatriptan* including:

- cold and flu medicines containing dextromethorphan (e.g. Robitussin Dry Cough Forte®)

Tell your pharmacist or doctor about all medicines or treatments that you may be taking, including vitamins, herbal products (e.g. St John's wort) or recreational drugs.

What side effects might you notice?

Side Effects	Recommended action
Pain, tightness or pressure in your chest, throat or jaw Fast or irregular heartbeat	Tell your doctor immediately
Drowsiness, tiredness or weakness, dizziness Hot flushing Burning sensation, tingling or numbness Pain, tenderness or redness at injection site	Tell your doctor if troublesome

If you notice any other effects, discuss them with your doctor or pharmacist.

Other information:

- Tell your doctor if you need to use *sumatriptan* several times each week. Using *sumatriptan* too often can cause a different type of headache. Your doctor can talk with you about different medicines that are taken every day to prevent migraines or cluster headaches.
- Do not use *sumatriptan* to prevent migraines or cluster headaches.
- *Sumatriptan* can impair your ability to do tasks such as driving or using machines. Alcohol makes this worse. Discuss your risk with your health professional. (search NZTA - Are you safe to drive?)